PROCEEDINGS OF SCIENCE

arXiv:1212.6823v2 [hep-ph] 2 Jan 2013

DESY 12-232, DO-TH-12/38, SFB/CPP-12-106, LPN12-142 `arXive:1212.6823[hep-ph]`

Three-Loop Contributions to the Gluonic Massive Operator Matrix Elements at General Values of N

Jakob Ablinger[a,b], **Johannes Blümlein**[*b], **Abilio De Freitas**[a], **Alexander Hasselhuhn**[a,b], **Sebastian Klein**[c], **Clemens Raab**[b], **Mark Round**[a], **Carsten Schneider**[a], **Fabian Wißbrock**[a]

[a]*Research Institute for Symbolic Computation (RISC), Johannes Kepler University Linz, Altenberger Str. 69, A-4040 Linz, Austria*

[b]*Deutsches Elektronen–Synchrotron (DESY), Zeuthen Platanenallee 6, D-15738 Zeuthen, Germany*

[c] *Institut für Theoretische Physik E, RWTH Aachen University, D-52056 Aachen, Germany*

Recent results on the calculation of 3-loop massive operator matrix elements in case of one and two heavy quark masses are reported. They concern the $O(n_f T_F^2 C_{F,A})$ and $O(T_F^2 C_{F,A})$ gluonic corrections, two-mass quarkonic moments, and ladder- and Benz-topologies. We also discuss technical aspects of the calculations.

Loops and Legs in Quantum Field Theory - 11th DESY Workshop on Elementary Particle Physics,
April 15-20, 2012
Wernigerode, Germany

---

[*] Speaker.

© Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike Licence.

1. Introduction

One of the most precise determinations of the strong coupling constant $\alpha_s(M_Z)$ relies on the next-to-next-to-leading order (NNLO) QCD analysis of the world deep-inelastic data [1,2]. Here currently the heavy flavor corrections to the structure function $F_2(x,Q^2)$ and transversity are known for the first Mellin moments $N = 2,..,10(14)$ [3,4].[1] In the asymptotic region $Q^2 \gg m^2$ the heavy flavor Wilson coefficients can be represented in form of convolutions [7] of massive operator matrix elements (OMEs) and the massless Wilson coefficients [8]. In case of the charm quark contribution the corresponding region is given by $Q^2/m_c^2 \gtrsim 10$. To carry out complete NNLO QCD analyses in this region the heavy flavor Wilson coefficients have to be known for general values of N. This also applies to precision measurements of the charm quark mass using the world deep-inelastic data [9].

Since 2010 the systematic calculation of the asymptotic massive Wilson coefficients at general values of N have been carried out. Five massive Wilson coefficients contribute to $F_2(x,Q^2)$ at NNLO [3]. Furthermore, there are other massive OMEs needed to compute the matching coefficients in the variable flavor number scheme in which the heavy quarks are assumed to decouple singly [3,10,11]. There are yet other contributions at NNLO from graphs containing both a massive charm and a bottom quark line, which extend the former representation in Ref. [3] and are necessary because of the fact that charm quarks do not yet become massless at the mass scale of bottom quarks since $m_c^2/m_b^2 \approx 1/10$ only. Results on first moments for these contributions are obtained in [12–14]. The extension of the renormalization of the massive OMEs is given in [14]. The logarithmic contributions at general values of N for the contributions to $F_2(x,Q^2)$ are available [15–17].[2] Furthermore, the NNLO heavy flavor Wilson coefficients in the asymptotic region were calculated for the structure function $F_L(x,Q^2)$ [3,20].

Two of the five massive Wilson coefficients contributing to $F_2(x,Q^2)$ at NNLO are known completely [21]. They are of $O(n_f T_F^2 C_{F,A})$. Likewise, these contributions to the further three Wilson coefficients and transversity were calculated for these color coefficients in [21]. Also the complete contribution $O(T_F^2 C_{A,F})$ for the OMEs $A_{Qq}^{\rm PS}, A_{qq,Q}^{\rm NS}, A_{qq,Q}^{\rm TR}$ are available [12].

In these proceedings we report on the calculation of the gluonic OMEs $O(n_f T_F^2 C_{F,A})$ [22] in Section 2 and on first results in $O(T_F^2 C_{F,A})$ for this channel in Section 3. Furthermore, the scalar 3-loop integrals for all ladder type integrals were calculated [23,24]. An extension of the method [25] to calculate finite Feynman integrals to the case of massive quark lines with local operator insertions has been used to calculate ladder- and Benz-topologies [23,24], also leading to new types of finite nested sums extending the harmonic [26], generalized harmonic [27,28], and cyclotomic sums [29] and the associated polylogarithms, cf. Section 4. Section 5 contains the conclusions.

2. Gluonic OMEs $\mathbf{O(n_f T_F^2 C_{F,A})}$

The calculation of all $O(n_f T_F^2 C_{F,A})$ contributions to the massive OMEs has been completed with the computation of $A_{gg,Q}$ and $A_{gq,Q}$ at this order in [22]. In these and other computations described below we used the codes `QGRAF` [30], `Form` and `tform` [31], and `color` [32]. For the check of individual moments of the expressions derived we also used `MATAD` [33].

---

[1] Present analyses use the NLO corrections in x-space [5] or Mellin space implementations [6].

[2] The $O(\alpha_s^2)$ and $O(\alpha_s^2 \varepsilon)$ contributions, with $\varepsilon = D - 4$ the dimensional parameter, needed in the renormalization were given in [7,18,19].

In the calculation of the $O(n_f T_F^2 C_{F,A})$-terms large amounts of nested sums emerge. An important step in the calculation consists in merging these individual sums to a smaller amount of sums containing very voluminous summands which are then solved with the codes used within the package `Sigma` [34], like `EvaluateMultiSum` and `SumProduction`, [35], written in `mathematica` [36], cf. also [37]. Moreover, this compactification shall be performed for whole diagrams to avoid the intermediate emergence of a larger amount of generalized harmonic sums as was observed in [21]. Generalized harmonic sums do not contribute in this case.

The constant contributions to the unrenormalized massive OMEs $A_{gg,Q}$ and $A_{gq,Q}$, $a_{gj,Q}^{(3)}$, $j = q, g$, read :

$$
\begin{aligned}
a_{gq,Q}^{(3),n_f T_F^2} = {} & C_F T_F^2 n_f \Biggl\{ -\frac{16\left(N^2+N+2\right)}{9(N-1)N(N+1)} \left(\frac{1}{3}S_1^3 + S_2 S_1 + \frac{2}{3}S_3 + 14\zeta_3 + 3S_1\zeta_2\right) \\
& + \frac{16\left(8N^3+13N^2+27N+16\right)}{27(N-1)N(N+1)^2}\left(3\zeta_2 + S_1^2 + S_2\right) \\
& - \frac{32\left(35N^4+97N^3+178N^2+180N+70\right)}{27(N-1)N(N+1)^3} S_1 \\
& + \frac{32\left(1138N^5+4237N^4+8861N^3+11668N^2+8236N+2276\right)}{243(N-1)N(N+1)^4}\Biggr\}, \qquad (2.1)
\end{aligned}
$$

$$
\begin{aligned}
a_{gg,Q}^{(3),n_f T_F^2} = {} & n_f T_F^2 \Biggl\{ C_A \frac{1}{(N-1)(N+2)} \Biggl[\frac{4P_1}{27N^2(N+1)^2} S_1^2 + \frac{8P_2}{729N^3(N+1)^3} S_1 \\
& + \frac{160}{27}(N-1)(N+2)\zeta_2 S_1 - \frac{448}{27}(N-1)(N+2)\zeta_3 S_1 + \frac{P_3}{729N^4(N+1)^4} \\
& - \frac{2P_4}{27N^2(N+1)^2}\zeta_2 + \frac{56\left(3N^4+6N^3+13N^2+10N+16\right)}{27N(N+1)}\zeta_3 - \frac{4P_5}{27N^2(N+1)^2} S_2 \Biggr] \\
& + C_F \frac{1}{(N-1)(N+2)} \Biggl[\frac{112\left(N^2+N+2\right)^2}{27N^2(N+1)^2} S_1^3 - \frac{16P_6}{27N^3(N+1)^3} S_1^2 \\
& + \frac{32P_7}{81N^4(N+1)^4} S_1 + \frac{16\left(N^2+N+2\right)^2}{3N^2(N+1)^2}\zeta_2 S_1 + \frac{16\left(N^2+N+2\right)^2}{3N^2(N+1)^2} S_2 S_1 \\
& - \frac{32P_8}{243N^5(N+1)^5} - \frac{16P_9}{9N^3(N+1)^3}\zeta_2 + \frac{448\left(N^2+N+2\right)^2}{9N^2(N+1)^2}\zeta_3 + \frac{16P_{10}}{9N^3(N+1)^3} S_2 \\
& - \frac{160\left(N^2+N+2\right)^2}{27N^2(N+1)^2} S_3 \Biggr]\Biggr\}. \qquad (2.2)
\end{aligned}
$$

The corresponding $1/\varepsilon$ terms contain contributions of the 3-loop anomalous dimensions [38] which we have verified by an independent calculation. Moreover a prediction made in [39] has been confirmed.

The gluonic OMEs $A_{gg,Q}$ and $A_{gq,Q}$ are needed for correct flavor matching in case of the transition of a *single* heavy quark becoming light, cf. [3, 10]. Here the correct choice of the matching scale is of importance [40].

In extending the above calculation the computation of some topological classes of diagrams has been automated mapping the graph to expressions involving hypergeometric ${}_{p+1}F_p$-functions. The ε-expansion leads to nested sums being calculated using the package `Sigma` [34]. The automation will cover other classes soon, requiring more involved ways to match the initial functions.

3. Gluonic OMEs $\mathbf{O(T_F^2 C_{F,A})}$ and OMEs with massive fermion lines of two different masses

All basic scalar topologies contributing to the $O(T_F^2 C_{F,A})$ of $A_{gg,Q}$ and $A_{gq,Q}$ have been calculated. An example is given in (3.1) for the graph containing two massive triangles with $m_1 = m_2$,

$$
\begin{aligned}
I_{D2}(N,\varepsilon) \;=\; (-1)^N \Biggl\{ & -\frac{1}{12N\varepsilon} + \frac{\bigl(27N^2-5N+16\bigr)}{1440N(N+1)(2N-1)4^N}\binom{2N}{N}\left[\sum_{\mathrm{i_1=1}}^{N} \frac{4^{i_1}}{i_1^2}\frac{S_1(i_1-1)}{\binom{2i_1}{i_1}} - 7\zeta_3\right] \\
& -\frac{(S_{2,1}-S_3-7\zeta_3)}{90N} - \frac{\bigl(S_1^2-S_2\bigr)}{90(N-1)N^2(N+1)} \\
& +\frac{\bigl(60N^3-19N^2-85N+60\bigr)}{720(N-1)N(N+1)(2N-1)}S_1 + \frac{\bigl(-162N^3+281N^2-187N+30\bigr)}{720(N-1)N^2(2N-1)}\Biggr\}. \qquad (3.1)
\end{aligned}
$$

Here we chose a minimal representation of sums being pairwise transcendent, which is proven by `Sigma` [34]. In these and similar diagrams sums of the kind

$$
\sum_{i=1}^{N} \frac{4^i}{i^3}\frac{1}{\binom{2i}{i}} \;=\; \int_0^1 dx \frac{x^N-1}{x-1}\mathrm{H}^*_{\mathsf{0w_3}}(x) \qquad (3.2)
$$

$$
\sum_{i=1}^{N} \frac{4^i}{i^2}\frac{1}{\binom{2i}{i}}S_1(i) \;=\; \int_0^1 dx \frac{x^N-1}{x-1}\left[\mathrm{H}^*_{\mathsf{0w_3}}(x) - \mathrm{H}^*_{\mathsf{w_30}}(x) - \mathrm{H}^*_{\mathsf{w_31}}(x) - 2\ln(2)\mathrm{H}^*_{\mathsf{w_3}}(x)\right] \qquad (3.3)
$$

emerge, which can be represented as Mellin transforms of iterated integrals, extending the usual harmonic polylogarithms (HPLs) [41], where $\mathrm{H}^*_{\emptyset}(x) = 1, \mathrm{H}^*_{\mathsf{b},\tilde{\mathsf{c}}}(x) = \int_x^1 dy\, f_{\mathsf{b}}(y)\mathrm{H}^*_{\tilde{\mathsf{c}}}(y), H^*_1(x) = \ln(1-x)$, and $f_{\mathsf{w_3}}(x) = 1/(x\sqrt{1-x})$, cf. [42]. The relative transcendence of the respective HPLs is proven using differential field methods [43], cf. also [44]. In exceptional cases they may be obtained in terms of HPLs with root arguments. Eq. (3.1) obviously is recurrent in N. The asymptotic expansions of (3.2, 3.3) are given by

$$
\sum_{i=1}^{N} \frac{4^i}{i^3}\frac{1}{\binom{2i}{i}} \;=\; 6\zeta_2\ln(2) - \frac{7}{2}\zeta_3 - \sqrt{\frac{\pi}{N}}\left\{\frac{2}{3}\frac{1}{N} - \frac{9}{20}\frac{1}{N^2} + \frac{199}{1344}\frac{1}{N^3} + O\left(\frac{1}{N^4}\right)\right\} \qquad (3.4)
$$

$$
\begin{aligned}
\sum_{i=1}^{N} \frac{4^i}{i^2}\frac{1}{\binom{2i}{i}}S_1(i) \;=\; & 6\zeta_2\ln(2) + \frac{7}{2}\zeta_3 + \sqrt{\frac{\pi}{N}}\left\{4 + \frac{7}{18}\frac{1}{N} - \frac{817}{2400}\frac{1}{N^2} + \frac{3835}{37632}\frac{1}{N^3} + O\left(\frac{1}{N^4}\right)\right\} \\
& -\frac{\ln(\overline{N})}{\sqrt{N}}\left\{2 - \frac{5}{12}\frac{1}{N} + \frac{21}{320}\frac{1}{N^2} + \frac{223}{10752}\frac{1}{N^3} + O\left(\frac{1}{N^4}\right)\right\}, \qquad (3.5)
\end{aligned}
$$

with $\overline{N} = Ne^{\gamma_E}$ and γ_E the Euler-Mascheroni number. The poles in the complex plane are situated at non-positive integers and half-integers and (3.1) is thus a meromorphic function. The half-integer poles emerge algebraically and from structures like

$$\binom{2N}{N}\frac{1}{4^N} = \frac{\Gamma(N+1/2)}{\sqrt{\pi}\Gamma(N+1)} . \tag{3.6}$$

With these properties it can be defined in the complex N-plane. By similar arguments the analytic continuation for the whole $T_F^2 C_{F,A}$-contribution to $A_{gg,Q}$ is obtained.

First results on massive OMEs with two fermion lines for $m_1 \neq m_2$ have been reported in [12, 13] for the moments $N = 2, 4, 6$ for A_{Qg}. The calculation has been performed by mapping the OMEs to tadpoles which were computed using `qexp` [45]. The renormalization of these matrix elements generalizes the case of n_f massless and one massive fermion [3] and is given in Ref. [14]. Since $m_c^2/m_b^2 \sim 1/10$, charm cannot be treated as massless at the scale m_b^2, which makes the use of the variable flavor scheme, see [3], built on this assumption, very problematic in case of these contributions. On the other hand, the fixed flavor number scheme which can be used in precision deep-inelastic world data analyses, cf. [1], can naturally accommodate these terms. The corresponding calculation in case of general values of N is underway [46].

4. Ladder and Benz Topologies

In the following we discuss higher topologies which contribute to the massive OMEs A_{ij} at 3–loop order. These are ladder-, Benz-, and crossed box topologies with the respective local operator insertions (of up to four lines). The basic ladder topologies have been calculated in Ref. [23] up to 3-leg operator insertions. Here we consider the case of only one massive fermion line. As has been described in [47] the Feynman diagrams can be represented as multiply nested sums. The ladder diagrams with six massive propagators have representations in terms of the Appell function F_1 [48]. Most of the integrals can be solved using `Sigma` [34], including the pole structure in ε. This is presently more involved in case of operator insertions with more than two legs at six massive lines.

For the non-divergent graphs the extension of the method [25] to local operators and massive lines allows the calculation. We consider the diagram shown in Fig. 1.



Figure 1: 3-loop ladder diagram containing a 3-vertex local operator insertion.

The local operator insertion can be resummed introducing a subsidiary parameter x

$$\sum_{j=0}^{N} T_{4a}^{N-j} T_{4b}^{j} \to \sum_{N=0}^{\infty}\sum_{j=0}^{N} x^N T_{4a}^{N-j} T_{4b}^{j} = \sum_{N=0}^{\infty} \frac{(T_{4a}x)^N - (T_{4b}x)^N}{T_{4a} - T_{4b}}$$
$$= \frac{1}{T_{4a} - T_{4b}}\left[\frac{1}{1 - xT_{4a}} - \frac{1}{1 - xT_{4b}}\right] = \frac{x}{(1 - xT_{4a})(1 - xT_{4b})} . \tag{4.1}$$

The integral may then be performed using the method [25] and is expressed in terms of hyperlogarithms $\mathrm{L}_{\vec{a}}(x)$. They obey the relations

$$\mathrm{L}_{b,\vec{a}}(x) = \int_0^x \frac{dy}{y-b}\mathrm{L}_{\vec{a}}(y), \quad \mathrm{L}_{\emptyset}(y) = 1; \quad \mathrm{L}_{\underbrace{0,\dots,0}_{n}}(x) = \frac{1}{n!}\ln^n(x), \quad a_i, b \in \mathbb{R}\,. \tag{4.2}$$

As an intermediary result one obtains

$$\begin{aligned}
\tilde{I}_4(x) \;=\; & \Bigg[-\frac{1+x}{x^3}\mathrm{L}_{-1} - \frac{2x-1}{x^3}\mathrm{L}_{1/2} - \frac{3(1-x)}{x^3}\mathrm{L}_{1} - \frac{1-2x+x^2}{(1-x)x^3}\mathrm{L}_{0,-1} + \frac{1-2x^2}{x^3}\mathrm{L}_{0,1/2} \\
& -\frac{3-4x-3x^2+3x^3}{(1-x)x^3}\mathrm{L}_{0,1} - \frac{1-2x^2}{x^3}\mathrm{L}_{1,1/2} + \frac{(1-x)(2+3x)}{x^3}\mathrm{L}_{1,1}\Bigg]\zeta_3 \\
& +\frac{(1+x)}{2x^3}\left(3\mathrm{L}_{-1,0,0,1} - 2\mathrm{L}_{-1,0,1,1} - 3\mathrm{L}_{1,0,0,1}\right) + \frac{1}{x^2}\left(6\mathrm{L}_{0,0,1,1} - 4\mathrm{L}_{0,1,0,1} - \mathrm{L}_{0,1,1,1}\right) \\
& -\frac{(-1+2x)}{2x^3}\left[3\mathrm{L}_{1/2,0,0,1} - \mathrm{L}_{1/2,0,1,1} - 3\mathrm{L}_{1/2,1,0,1} + \mathrm{L}_{1/2,1,1,1}\right] \\
& -\frac{3}{2x^2}\mathrm{L}_{1,0,1,1} + \frac{2}{x^2}\mathrm{L}_{1,1,0,1} - \frac{(-1+x)}{2x^3}\mathrm{L}_{1,1,1,1} + \frac{2}{x^2}\left(\mathrm{L}_{0,1,1} - \mathrm{L}_{1,0,1}\right) \\
& +\frac{\left(-1+2x+x^2\right)}{2(-1+x)x^3}\left[3\mathrm{L}_{0,-1,0,0,1} - 2\mathrm{L}_{0,-1,0,1,1}\right] \\
& -\frac{5}{-1+x}\mathrm{L}_{0,0,0,1,1} - \frac{5}{2(-1+x)}\mathrm{L}_{0,0,1,0,1} + \frac{3(3+x)}{2(-1+x)x}\mathrm{L}_{0,0,1,1,1} \\
& -\frac{\left(-1+2x^2\right)}{2x^3}\left[3\mathrm{L}_{0,1/2,0,0,1} + \mathrm{L}_{0,1/2,0,1,1} + 3\mathrm{L}_{0,1/2,1,0,1} - \mathrm{L}_{0,1/2,1,1,1}\right] \\
& +\frac{3\left(1-3x^2+3x^3\right)}{2(-1+x)x^3}\mathrm{L}_{0,1,0,0,1} + \frac{8-14x+5x^2+3x^3}{2(-1+x)x^3}\mathrm{L}_{0,1,0,1,1} \\
& +\frac{8-15x+3x^2}{2(-1+x)x^3}\mathrm{L}_{0,1,1,0,1} - \frac{3(-3+2x)}{2x^3}\mathrm{L}_{0,1,1,1,1} + \frac{-6+3x+5x^2}{x^3}\mathrm{L}_{1,0,0,1,1} \\
& +\frac{2(-1+x)}{x^3}\mathrm{L}_{1,1,1,0,1} + \frac{4-2x+5x^2}{2x^3}\mathrm{L}_{1,0,1,0,1} - \frac{-4+6x+3x^2}{2x^3}\mathrm{L}_{1,0,1,1,1} \\
& +\frac{\left(-1+2x^2\right)}{2x^3}\left[3\mathrm{L}_{1,1/2,0,0,1} - \mathrm{L}_{1,1/2,0,1,1}\right] - \frac{3(-1+x)(4+3x)}{2x^3}\mathrm{L}_{1,1,0,0,1} \\
& -\frac{\left(-1+2x^2\right)}{2x^3}\left[\mathrm{L}_{1,1/2,1,0,1} + \mathrm{L}_{1,1/2,1,1,1}\right] - \frac{(-1+x)(5+3x)}{2x^3}\mathrm{L}_{1,1,0,1,1}\,.
\end{aligned} \tag{4.3}$$

Here iterated integrals over the alphabet $\{0,1,-1,1/2\}$ contribute. Now the Nth Taylor coefficient has to be obtained for $\tilde{I}_4(x)$, which is possible using the package `HarmonicSums` [28] :

$$\begin{aligned}
\hat{I}_4(N) \;=\; & \frac{P_1}{2(1+N)^5(2+N)^5(3+N)^5} + \frac{P_2}{(1+N)^2(2+N)^2(3+N)^2}\zeta_3 \\
& +\frac{(-1)^N\left(65+101N+56N^2+13N^3+N^4\right)}{2(1+N)^2(2+N)^2(3+N)^2}S_{-3} + \frac{\left(-24-5N+2N^2\right)}{12(2+N)^2(3+N)^2}S_1^3 \\
& -\frac{1}{2(1+N)(2+N)(3+N)}S_2^2 + \frac{1}{(2+N)(3+N)}S_1^2 S_2 \\
& +\frac{314+631N+578N^2+288N^3+68N^4+5N^5}{4(1+N)^3(2+N)^2(3+N)^2}S_1^2 - \frac{3}{2}S_5
\end{aligned}$$

$$
\begin{aligned}
&-\frac{\left(399+2069N+2774N^2+1510N^3+349N^4+27N^5\right)}{6(1+N)^2(2+N)^2(3+N)^2}S_3-2S_{-2,-3}\\
&-2\zeta_3S_{-2}-S_{-2,1}S_{-2}+\frac{(-1)^N\left(65+101N+56N^2+13N^3+N^4\right)}{(1+N)^2(2+N)^2(3+N)^2}S_{-2,1}\\
&+\frac{\left(59+42N+6N^2\right)}{2(1+N)(2+N)(3+N)}S_4+\frac{(5+N)}{(1+N)(3+N)}\zeta_3S_1\left(2\right)\\
&-\frac{752+2087N+2490N^2+1580N^3+558N^4+105N^5+8N^6}{4(1+N)^3(2+N)^2(3+N)^2}S_2-\zeta_3S_2\\
&-\frac{3}{2}S_3S_2-2S_{2,1}S_2+\frac{\left(99+225N+190N^2+65N^3+7N^4\right)}{2(1+N)^2(2+N)^2(3+N)}S_{2,1}\\
&+\frac{P_3}{(1+N)^4(2+N)^4(3+N)^4}S_1-\frac{(11+5N)}{(1+N)(2+N)(3+N)}\zeta_3S_1\\
&-\frac{\left(470+1075N+996N^2+447N^3+96N^4+8N^5\right)}{4(1+N)^2(2+N)^2(3+N)^2}S_2S_1-S_{2,3}\\
&+\frac{(53+29N)}{2(1+N)(2+N)(3+N)}S_3S_1-\frac{3(3+2N)}{(1+N)(2+N)(3+N)}S_1S_{2,1}\\
&+\frac{\left(-79-40N+N^2\right)}{2(1+N)(2+N)(3+N)}S_{3,1}-3S_{4,1}+S_{-2,1,-2}\\
&+\frac{2^{1+N}\left(-28-25N-4N^2+N^3\right)}{(1+N)^2(2+N)(3+N)^2}S_{1,2}\left(\frac{1}{2},1\right)-\frac{\left(-7+2N^2\right)}{(1+N)(2+N)(3+N)}S_{2,1,1}\\
&+5S_{2,2,1}+6S_{3,1,1}+\frac{2^N\left(-28-25N-4N^2+N^3\right)}{(1+N)^2(2+N)(3+N)^2}S_{1,1,1}\left(\frac{1}{2},1,1\right)\\
&-\frac{(5+N)}{(1+N)(3+N)}S_{1,1,2}\left(2,\frac{1}{2},1\right)-\frac{(5+N)}{2(1+N)(3+N)}S_{1,1,1,1}\left(2,\frac{1}{2},1,1\right)\,. \qquad (4.4)
\end{aligned}
$$

Here P_i denote polynomials, cf. [23]. The final expression contains individual terms which grow $\propto 2^N$ for $N\rightarrow\infty$. However, this singularity cancels in $\hat{I}_4(N)$. As an extension of the usual harmonic sums [26] also generalized sums [27,28] contribute with weights $x_i\in\{1,1/2,2\}$.

Let us now turn to the graph shown in Fig. 2. It consists out of two contributions with respect to the operator insertion, cf. [3], which may be viewed as being obtained by contraction of the central lines of the ladder graph, resp. the graph of the crossed box, with central operator insertion.



Figure 2: 3-loop ladder diagram containing a 4-vertex local operator insertion.

While the calculation of the former graph is rather straightforward, in the latter case root-letters appear in the integration formalism using the method of hyperlogarithms. Here it is possible, however, to move the corresponding root-expressions completely into the argument of the hyperlogarithms.

In the next step we would like to determine the Nth Taylor coefficient of the expression obtained. A first possibility consists in calculating a large number of Mellin-moments for the expression in an efficient way, which needs the use of `Form` [31] beyond the representations in `Maple` [49]. In the present case about 1500 moments have been calculated. The method of guessing [50] allows to derive a corresponding difference equation, for which we needed $\sim$ 700 moments in the present case. This equation can now be solved using `Sigma` [34] and the Nth moment is obtained. In an earlier investigation we have determined all 3-loop anomalous dimensions and massless Wilson coefficients from their moments in this way, which required 5114 moments, [51]. Although not expected to fail with a significant probability, and having a large verification space with $\sim$ 1500 moments available at $\sim$ 700 needed, still the Nth moment shall be derived having a proof certificate. A more involved calculation using `HarmonicSums` [28] and `Sigma` [34] provides this. Here, difference equations of up to order `o = 16` and degree `d = 108` have to be solved. The results are presented in [24].

The emergence of root-expressions in x implies in the present result quite a series of new sums, which are of the nested binomial- and inverse-binomial type and also contain generalized harmonic sums. We have translated the result into x-space, where new iterated integrals with various root-type letters were obtained, extending those having appeared in [52]. Using the methods of Ref. [43] the relative transcendence of the different functions can be checked and we derived a corresponding basis for these functions, cf. [42]. In N-space the basis representation, including the new sums, is derived using `Sigma`, [34].

The method of hyperlogarithms also allows the calculation of non-singular Benz-graphs, which have representations in terms of harmonic sums and generalized harmonic sums, cf. [24, 53].

5. Conclusions

We reported on recent progress in calculating the asymptotic heavy flavor Wilson coefficients contributing to the deep-inelastic structure function $F_2(x,Q^2)$ at 3-loop order. As a first class all contributions of $O(n_f T_F^2 C_{A,F})$ have been completed and basis integrals for the class $O(T_F^2 C_{A,F})$ were obtained. Furthermore, the automatic calculation of related topology classes for other color factors started. We extended the former analysis to the case in which two heavy quarks of different mass contribute and obtained a series of Mellin moments. These contributions are no longer in accordance with the variable flavor number scheme, since charm does not decouple at the mass scale of the bottom quark. In case of two massive fermion lines with equal mass new sums and iterated integrals appear beyond the usual harmonic sums and polylogarithms, also leading to more singularities in the complex plane. We have also calculated the scalar graphs contributing to ladder-topologies for up to six massive fermion lines, including graphs with local 4-leg operators. Here the results are given in terms of special classes of generalized sums, which individually may even diverge exponentially for $N \to \infty$. This divergence is canceled between different terms contributing. In case of the graph shown in Fig. 2 a larger amount of nested binomial and inverse binomial sums weighted with harmonic sums and their generalization, also including cyclotomic sums, emerge, extending the known alphabets both for the sums and the associated iterated integrals. This all is invisible at the level of Mellin moments since the corresponding expressions are given in terms of rationals and single ζ-values. It appears that in the presence of a single mass already at 3-loop order rich new structures are contributing in the single differential case being characterized either by the Mellin variable N or the momentum fraction x.

Acknowledgment. We would like to thank F. Brown for discussions. The Feynman diagrams have been drawn using `Axodraw` [54]. This work has been supported in part by DFG Sonderforschungsbereich Transregio 9, Computergestützte Theoretische Teilchenphysik, by the Austrian Science Fund (FWF) grant P203477-N18, by the EU Network LHCPHENOnet PITN-GA-2010-264564, and ERC Starting Grant PAGAP FP7-257638. We thank WolframResearch for their kind cooperation within LHCPHENOnet.

References

[1] S. Alekhin, J. Blümlein and S. Moch, *Parton Distribution Functions and Benchmark Cross Sections at NNLO*, Phys. Rev. D **86** (2012) 054009 [arXiv:1202.2281 [hep-ph]].

[2] S. Bethke *et al.*, *Workshop on Precision Measurements of α_s*, arXiv:1110.0016 [hep-ph].

[3] I. Bierenbaum, J. Blümlein and S. Klein, *Mellin Moments of the $O(\alpha_s^3)$ Heavy Flavor Contributions to unpolarized Deep-Inelastic Scattering at $Q^2 \gg m^2$ and Anomalous Dimensions*, Nucl. Phys. B **820** (2009) 417, [arXiv:0904.3563 [hep-ph]].

[4] J. Blümlein, S. Klein and B. Tödtli, *$O(\alpha_s^2)$ and $O(\alpha_s^3)$ Heavy Flavor Contributions to Transversity at $Q^2 \gg m^2$*, Phys. Rev. D **80** (2009) 094010 [arXiv:0909.1547 [hep-ph]].

[5] E. Laenen, S. Riemersma, J. Smith and W. L. van Neerven, *Complete $O(\alpha_s)$ corrections to heavy flavor structure functions in electroproduction* Nucl. Phys. B **392** (1993) 162; 229;
S. Riemersma, J. Smith and W. L. van Neerven, *Rates for inclusive deep inelastic electroproduction of charm quarks at HERA*, Phys. Lett. B **347** (1995) 143 [hep-ph/9411431].

[6] S. I. Alekhin and J. Blümlein, *Mellin representation for the heavy flavor contributions to deep inelastic structure functions*, Phys. Lett. B **594** (2004) 299 [hep-ph/0404034].

[7] M. Buza, Y. Matiounine, J. Smith, R. Migneron and W. L. van Neerven, *Heavy quark coefficient functions at asymptotic values $Q^2 \gg m^2$*, Nucl. Phys. B **472** (1996) 611 [hep-ph/9601302].

[8] J. A. M. Vermaseren, A. Vogt and S. Moch, *The Third-order QCD corrections to deep-inelastic scattering by photon exchange*, Nucl. Phys. B **724** (2005) 3 [hep-ph/0504242].

[9] S. Alekhin, J. Blümlein, K. Daum, K. Lipka and S. Moch, *Precise charm-quark mass from deep-inelastic scattering*, arXiv:1212.2355 [hep-ph].

[10] M. Buza, Y. Matiounine, J. Smith and W. L. van Neerven, *Charm electroproduction viewed in the variable flavor number scheme versus fixed order perturbation theory*, Eur. Phys. J. C **1** (1998) 301 [hep-ph/9612398].

[11] I. Bierenbaum, J. Blümlein and S. Klein, *The Gluonic Operator Matrix Elements at $O(\alpha_s^2)$ for DIS Heavy Flavor Production*, Phys. Lett. B **672** (2009) 401 [arXiv:0901.0669 [hep-ph]].

[12] J. Ablinger, J. Blümlein, S. Klein, C. Schneider and F. Wißbrock, *3-Loop Heavy Flavor Corrections to DIS with two Massive Fermion Lines*, arXiv:1106.5937 [hep-ph].

[13] J. Ablinger, J. Blümlein, A. Hasselhuhn, S. Klein, C. Schneider and F. Wißbrock, *New Heavy Flavor Contributions to the DIS Structure Function $F_2(x,Q^2)$ at $O(\alpha_s^3)$*, PoS (RADCOR2011) 031 arXiv:1202.2700 [hep-ph].

[14] J. Blümlein and F. Wißbrock, in preparation.

[15] I. Bierenbaum, J. Blümlein and S. Klein, *Logarithmic $O(\alpha_s^3)$ contributions to the DIS Heavy Flavor Wilson Coefficients at $Q^2 \gg m^2$*, PoS DIS **2010** (2010) 148 [arXiv:1008.0792 [hep-ph]].

[16] I. Bierenbaum, J. Blümlein, S. Klein, and F. Wißbrock, in preparation.

[17] H. Kawamura, N. A. Lo Presti, S. Moch and A. Vogt, *On the next-to-next-to-leading order QCD corrections to heavy-quark production in deep-inelastic scattering*, Nucl. Phys. B **864** (2012) 399 [arXiv:1205.5727 [hep-ph]].

[18] I. Bierenbaum, J. Blümlein and S. Klein, *Two-Loop Massive Operator Matrix Elements and Unpolarized Heavy Flavor Production at Asymptotic Values $Q^2 \gg m^2$*, Nucl. Phys. B **780** (2007) 40 [hep-ph/0703285].

[19] I. Bierenbaum, J. Blümlein, S. Klein and C. Schneider, *Two-Loop Massive Operator Matrix Elements for Unpolarized Heavy Flavor Production to $O(\varepsilon)$*, Nucl. Phys. B **803** (2008) 1 [arXiv:0803.0273 [hep-ph]].

[20] J. Blümlein, A. De Freitas, W. L. van Neerven and S. Klein, *The Longitudinal Heavy Quark Structure Function F_L^{QQ} in the Region $Q^2 \gg m^2$ at $O(\alpha_s^3)$*, Nucl. Phys. B **755** (2006) 272 [hep-ph/0608024].

[21] J. Ablinger, J. Blümlein, S. Klein, C. Schneider and F. Wißbrock, *The $O(\alpha_s^3)$ Massive Operator Matrix Elements of $O(N_f)$ for the Structure Function $F_2(x,Q^2)$ and Transversity*, Nucl. Phys. B **844** (2011) 26 [arXiv:1008.3347 [hep-ph]].

[22] J. Blümlein, A. Hasselhuhn, S. Klein and C. Schneider, *The $O(\alpha_s^3 n_f T_F^2 C_{A,F})$ Contributions to the Gluonic Massive Operator Matrix Elements*, Nucl. Phys. B **866** (2013) 196 [arXiv:1205.4184 [hep-ph]].

[23] J. Ablinger, J. Blümlein, A. Hasselhuhn, S. Klein, C. Schneider and F. Wißbrock, *Massive 3-loop Ladder Diagrams for Quarkonic Local Operator Matrix Elements*, Nucl. Phys. B **864** (2012) 52 [arXiv:1206.2252 [hep-ph]].

[24] J. Ablinger, J. Blümlein, C. Schneider, and F. Wißbrock, in prepration.

[25] F. Brown, *The Massless higher-loop two-point function*, Commun. Math. Phys. **287** (2009) 925 [arXiv:0804.1660 [math.AG]].

[26] J. A. M. Vermaseren, *Harmonic sums, Mellin transforms and integrals*, Int. J. Mod. Phys. A **14** (1999) 2037 [hep-ph/9806280];
J. Blümlein and S. Kurth, *Harmonic sums and Mellin transforms up to two loop order*, Phys. Rev. D **60** (1999) 014018 [hep-ph/9810241].

[27] S. Moch, P. Uwer and S. Weinzierl, *Nested sums, expansion of transcendental functions and multiscale multiloop integrals*, J. Math. Phys. **43** (2002) 3363 [hep-ph/0110083];

[28] J. Ablinger, J. Blümlein and C. Schneider, *Analytic and Algorithmic Aspects of Generalized Harmonic Sums and Polylogarithms*, DESY 12–210;
J. Ablinger, *Computer Algebra Algorithms for Special Functions in Particle Physics*, J. Kepler University Linz. PhD Thesis. April 2012;
`http://www.risc.jku.at/research/combinat/software/`

[29] J. Ablinger, J. Blümlein and C. Schneider, *Harmonic Sums and Polylogarithms Generated by Cyclotomic Polynomials*, J. Math. Phys. **52** (2011) 102301 [arXiv:1105.6063 [math-ph]].

[30] P. Nogueira, *Automatic Feynman graph generation*, J. Comput. Phys. **105** (1993) 279.

[31] J. A. M. Vermaseren, *New features of FORM*, math-ph/0010025;
M. Tentyukov and J. A. M. Vermaseren, *The Multithreaded version of FORM*, Comput. Phys. Commun. **181** (2010) 1419 [hep-ph/0702279 [HEP-PH]].

[32] T. van Ritbergen, A. N. Schellekens and J. A. M. Vermaseren, *Group theory factors for Feynman diagrams*, Int. J. Mod. Phys. A **14** (1999) 41 [arXiv:hep-ph/9802376].

[33] M. Steinhauser, *MATAD: A Program package for the computation of MAssive TADpoles*, Comput. Phys. Commun. **134** (2001) 335 [hep-ph/0009029].

[34] C. Schneider, *A refined difference field theory for symbolic summation* J. Symbolic Comput. **43** (2008) 611 [arXiv:0808.2543]; *Product representations in $\Pi\Sigma$-fields* Ann. Comb. **9** (2005) 75; *Solving parameterized linear difference equations in terms of indefinite nested sums and products*, J. Differ. Equations Appl. **11** (2005) 799; *Parameterized Telescoping Proves Algebraic Independence of Sums* Ann. Comb. **14** (2010) 533 [arXiv:0808.2596]; *A Symbolic Summation Approach to Find Optimal Nested Sum Representations*. In A. Carey, D. Ellwood, S. Paycha, and S. Rosenberg, editors, *Motives, Quantum Field Theory, and Pseudodifferential Operators*, volume 12 of *Clay Mathematics Proceedings* 285, Amer. Math. Soc, 2010 arXiv:0808.2543; *Symbolic summation assists combinatorics* Sém. Lothar. Combin., **56** (2007) 1, Article B56b; *Multisummation in Difference Fields*, Habilitationsschrift JKU Linz (2007) and references therein;
J. Ablinger, J. Blümlein, S. Klein and C. Schneider, *Modern Summation Methods and the Computation of 2- and 3-loop Feynman Diagrams*, Nucl. Phys. Proc. Suppl. **205-206** (2010) 110 [arXiv:1006.4797 [math-ph]].

[35] C. Schneider, in preparation.

[36] `http://www.wolfram.com/mathematica/`

[37] J. Blümlein, A. Hasselhuhn and C. Schneider, *Evaluation of Multi-Sums for Large Scale Problems*, arXiv:1202.4303 [math-ph]; in: Proceedings of 10th International Symposium on Radiative Corrections, PoS(RADCOR2011)32, (2012);
J. Ablinger, S. Blümlein, M. Round and C. Schneider, *Advanced Computer Algebra Algorithms for the Expansion of Feynman Integrals*, arXiv:1210.1685 [cs.SC].

[38] A. Vogt, S. Moch and J. A. M. Vermaseren, *The Three-loop splitting functions in QCD: The Singlet case*, Nucl. Phys. B **691** (2004) 129 [hep-ph/0404111].

[39] J. F. Bennett and J. A. Gracey, *Determination of the anomalous dimension of gluonic operators in deep inelastic scattering at $O(1/N_f)$*, Nucl. Phys. B **517** (1998) 241 [hep-ph/9710364].

[40] J. Blümlein and W. L. van Neerven, *Heavy flavor contributions to the deep inelastic scattering sum rules*, Phys. Lett. B **450** (1999) 417 [hep-ph/9811351].

[41] E. Remiddi and J. A. M. Vermaseren, *Harmonic polylogarithms*, Int. J. Mod. Phys. A **15** (2000) 725 [hep-ph/9905237].

[42] J. Ablinger, J. Blümlein, C. Raab and C. Schneider, in preparation.

[43] M. Bronstein, *Symbolic Integration I - Transcendental Functions*, (Springer, Heidelberg, 1997).

[44] C. Raab, *Definite Integration in Differential Fields*, PhD Tesis, August 2012, Johannes Kepler University Linz, Austria.

[45] R. Harlander, T. Seidensticker, M. Steinhauser, *Complete corrections of $O(\alpha\alpha_s)$ to the decay of the Z-boson into bottom quarks*, *Phys. Lett.* **B426** (1998) 125, [hep-ph/9712228];
T. Seidensticker, *Automatic application of successive asymptotic expansions of Feynman diagrams*, [hep-ph/9905298].

[46] J. Blümlein, M. Round, C. Schneider, F. Wißbrock, in preparation.

[47] J. Blümlein, *Structural Relations of Harmonic Sums and Mellin Transforms up to Weight w = 5*, Comput. Phys. Commun. **180** (2009) 2218 [arXiv:0901.3106 [hep-ph]];
J. Blümlein, S. Klein, C. Schneider and F. Stan, *A Symbolic Summation Approach to Feynman Integral Calculus*, J. Symbolic Comput. 47 (2012) 1267-1289 [arXiv:1011.2656 [cs.SC]].

[48] P. Appell, *Sur Les Fonctions Hypérgéometriques de Plusieurs Variables*, (Gauthier-Villars, Paris, 1925);
P. Appell and J. Kampé de Fériet, *Fonctions Hypérgéometriques; Polynômes d'Hermite*, (Gauthier-Villars, Paris, 1926);
W.N. Bailey, *Generalized Hypergeometric Series*, (Cambridge University Press, Cambridge, 1935);
A. Erdélyi (ed.), *Higher Transcendental Functions*, Bateman Manuscript Project, Vol. **I**, (McGraw-Hill, New York, 1953);
L.J. Slater, *Generalized Hypergeometric Functions*, (Cambridge University Press, Cambridge, 1966);
H. Exton, *Multiple Hypergeometric Functions and Applications*, (Ellis Horwood Limited, Chichester, 1976);
H. Exton, *Handbook of Hypergeometric Integrals*, (Ellis Horwood Limited, Chichester, 1978).

[49] `http://www.maplesoft.com/products/maple/`

[50] M. Kauers, *Guessing Handbook*, Technical Report RISC 09-07 (2009), JKU Linz.

[51] J. Blümlein, M. Kauers, S. Klein and C. Schneider, *Determining the closed forms of the* $O(\alpha_s^3)$ *anomalous dimensions and Wilson coefficients from Mellin moments by means of computer algebra*, Comput. Phys. Commun. **180** (2009) 2143 [arXiv:0902.4091 [hep-ph]].

[52] U. Aglietti and R. Bonciani, *Master integrals with 2 and 3 massive propagators for the 2 loop electroweak form-factor - planar case*, Nucl. Phys. B **698** (2004) 277 [hep-ph/0401193].

[53] J. Ablinger, J. Blümlein, A. De Freitas, A. Hasselhuhn, S. Klein, C. Schneider and F. Wißbrock, *New Results on the 3-Loop Heavy Flavor Wilson Coefficients in Deep-Inelastic Scattering*, arXiv:1212.5950 [hep-ph], PoS(ICHEP2012)270.

[54] J. A. M. Vermaseren, *Axodraw*, Comput. Phys. Commun. **83** (1994) 45.